

(19) World Intellectual Property Organization
International Bureau




(43) International Publication Date
3 January 2002 (03.01.2002)

PCT


(10) International Publication Number
WO 02/00209 A2

- (51) International Patent Classification⁷: **A61K 31/00**
- (21) International Application Number: **PCT/US01/16343**
- (22) International Filing Date: **18 May 2001 (18.05.2001)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
60/214,171 **26 June 2000 (26.06.2000)** **US**
- (71) Applicant (*for all designated States except US*):
WARNER-LAMBERT COMPANY [US/US]; 201 Tabor Road, Morris Plains, NJ 07950 (US).
- (74) Agents: **FEDERMAN, Evan, J. et al.;** Warner-Lambert Company, 201 Tabor Road, Morris Plains, NJ 07950 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **BRYANS, Justin, Stephen [GB/GB];** Pfizer Global Research & Development, Forvie Site, Robinson Way, Cambridge CB2 2QB, Cambridgeshire (GB). **MELTZER, Leonard, Theodore [US/US];** Pfizer Inc., PGRD, Ann Arbor Laboratories, 2800 Plymouth Road, Ann Arbor, MI 48105 (US).
- Published:
— *without international search report and to be republished upon receipt of that report*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: **GABAPENTIN ANALOGUES FOR SLEEP DISORDERS**



(1)



(1A)

(57) Abstract: The invention provides a new use of compounds for formula 1 or 1A or a pharmaceutically acceptable salt thereof. The compounds are useful in the treatment of insomnia and related disorders.

WO 02/00209 A2


US 10/735,561 Filed: 12/12/2003 Art Unit: 1614
Docket: PC25627A Inventor: Dooley, et al.

-1-

GABAPENTIN ANALOGUES FOR SLEEP DISORDERS

BACKGROUND OF THE INVENTION


Compounds of formula




5 wherein R₁ is hydrogen or a lower alkyl radical and n is 4, 5, or 6 are known in United States Patent Number 4,024,175 and its divisional United States Patent Number 4,087,544. The uses disclosed are: protective effect against cramp induced by thiosemicarbazide; protective action against cardiazole cramp; the cerebral diseases, epilepsy, faintness attacks, hypokinesia, and cranial traumas; and improvement in cerebral functions. The compounds are useful in geriatric patients. The patents are hereby incorporated by reference.

United States Patent Application 09/485,382 filed February 8, 2000, covers compounds of formulas 1 and 1A below. The application discloses various utilities for the compounds. This is incorporated by reference.

15 Compounds of Formula I



wherein R₁ is hydrogen or lower alkyl and n is an integer of from 4 to 6, and the pharmaceutically acceptable salts thereof and compounds of Formula II



-2-

hydrogen, methyl, or carboxyl; or an individual enantiomeric isomer thereof; or a pharmaceutically acceptable salt thereof, are useful in the treatment of insomnia (United States Patent Application 60/092166 filed July 9, 1998, which is incorporated here by reference).

5

SUMMARY OF THE INVENTION

The instant invention is a method of treating insomnia, that is, difficulty in sleeping or disturbed sleep patterns which leave the perception of insufficient sleep. Insomnia is a common symptom which may be due to several emotional and physical disorders (The Merck Manual, 16th ed., pp 1445-6).

10

The benefit of using compounds of the invention to treat insomnia is that they are not addictive. Additionally, they have a half-life in the body that is suitable to work during the evening and subsequently clear the body by morning to allow for easy arousal. The compounds can be combined with other agents to enhance the sleep inducing effects. Such agents include melatonin, tryptophan, valerian, passiflora, antihistamines such as diphenhydramine hydrochloride or doxylamine succinate, benzodiazepines, and nonbenzodiazepines hypnotics.


15

Additional advantages of using the compounds of formula 1 and 1A in the present invention include the relatively nontoxic nature of the compounds, the ease of preparation, the fact that the compounds are well-tolerated, and the ease of IV administration of the drugs. The subjects treated with the method of the present invention are mammals, including humans.

20


The compounds useful in the practice of the invention are those of formulas 1 and 1A

-3-



1

or




1A

wherein R to R¹⁴ are as defined below.

The compounds of the invention and their pharmaceutically acceptable salts and the prodrugs of the compounds are useful in the treatment of insomnia and sleeplessness.


DETAILED DESCRIPTION OF THE INVENTION

The compounds useful in the instant invention and their pharmaceutically acceptable salts are as defined by formulas 1 and 1A



1

and



1A

or a pharmaceutically acceptable salt thereof wherein:

R is hydrogen or a lower alkyl;

R¹ to R¹⁴ are each independently selected from hydrogen, straight or branched alkyl of from 1 to 6 carbons, phenyl, benzyl, fluorine, chlorine, bromine, hydroxy, hydroxymethyl, amino, aminomethyl, trifluoromethyl, -CO₂H, -CO₂R¹⁵, -CH₂CO₂H, -CH₂CO₂R¹⁵, -OR¹⁵ wherein R¹⁵ is a straight or branched alkyl of from 1 to 6 carbons, phenyl, or benzyl, and R¹ to R⁸ are not simultaneously hydrogen.

-4-

Preferred compounds of the invention are those of Formula I wherein R¹ to R¹⁴ are selected from hydrogen, methyl, ethyl, propyl, isopropyl, butyl straight or branched, phenyl, or benzyl.

5 More preferred compounds are those of Formula I wherein R¹ to R¹⁴ are selected from hydrogen, methyl, ethyl, or benzyl.

The most preferred compounds are selected from:

- (1 α ,3 α ,4 α)-(1-Aminomethyl-3,4-dimethyl-cyclopentyl)-acetic acid;
(1 α ,3 α ,4 α)-(1-Aminomethyl-3,4-diethyl-cyclopentyl)-acetic acid;
(1 α ,3 α ,4 α)-(1-Aminomethyl-3,4-diisopropyl-cyclopentyl)-acetic acid;
10 [1S-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-ethyl-4-methyl-cyclopentyl)-acetic acid;
[1R-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-ethyl-4-methyl-cyclopentyl)-acetic acid;
[1S-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-isopropyl-4-methyl-cyclopentyl)-
15 acetic acid;
[1R-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-isopropyl-4-methyl-cyclopentyl)-acetic acid;
[1S-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-ethyl-4-isopropyl-cyclopentyl)-acetic acid;
20 [1R-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-ethyl-4-isopropyl-cyclopentyl)-acetic acid;
[1S-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-tert-butyl-4-methyl-cyclopentyl)-acetic acid;
[1R-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-tert-butyl-4-methyl-cyclopentyl)-
25 acetic acid;
[1S-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-tert-butyl-4-ethyl-cyclopentyl)-acetic acid;
[1R-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-tert-butyl-4-ethyl-cyclopentyl)-acetic acid;

-5-

- [1S-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-tert-butyl-4-isopropyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-tert-butyl-4-isopropyl-cyclopentyl)-acetic acid;
- 5 (1 α ,3 α ,4 α)-(1-Aminomethyl-3,4-di-tert-butyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-methyl-4-phenyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-methyl-4-phenyl-cyclopentyl)-acetic acid;
- 10 [1S-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-benzyl-4-methyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-benzyl-4-methyl-cyclopentyl)-acetic acid;
- (1S-cis)-(1-Aminomethyl-3-methyl-cyclopentyl)-acetic acid;
- 15 (1S-cis)-(1-Aminomethyl-3-ethyl-cyclopentyl)-acetic acid;
- (1S-cis)-(1-Aminomethyl-3-isopropyl-cyclopentyl)-acetic acid;
- (1S-cis)-(1-Aminomethyl-3-tert-butyl-cyclopentyl)-acetic acid;
- (1S-cis)-(1-Aminomethyl-3-phenyl-cyclopentyl)-acetic acid;
- (1S-cis)-(1-Aminomethyl-3-benzyl-cyclopentyl)-acetic acid;
- 20 (1R-cis)-(1-Aminomethyl-3-methyl-cyclopentyl)-acetic acid;
- (1R-cis)-(1-Aminomethyl-3-ethyl-cyclopentyl)-acetic acid;
- (1R-cis)-(1-Aminomethyl-3-isopropyl-cyclopentyl)-acetic acid;
- (1R-cis)-(1-Aminomethyl-3-tert-butyl-cyclopentyl)-acetic acid;
- (1R-cis)-(1-Aminomethyl-3-phenyl-cyclopentyl)-acetic acid;
- 25 (1R-cis)-(1-Aminomethyl-3-benzyl-cyclopentyl)-acetic acid;
- (S)-(1-Aminomethyl-3,3-dimethyl-cyclopentyl)-acetic acid;
- (S)-(1-Aminomethyl-3,3-diethyl-cyclopentyl)-acetic acid;
- (1-Aminomethyl-3,3,4,4-tetramethyl-cyclopentyl)-acetic acid;
- (1-Aminomethyl-3,3,4,4-tetraethyl-cyclopentyl)-acetic acid;
- 30 (1 α ,3 β ,4 β)-(1-Aminomethyl-3,4-dimethyl-cyclopentyl)-acetic acid;
- (1 α ,3 β ,4 β)-(1-Aminomethyl-3,4-diethyl-cyclopentyl)-acetic acid;

-6-

- (1 α ,3 β ,4 β)-(1-Aminomethyl-3,4-diisopropyl-cyclopentyl)-acetic acid;
[1R-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-ethyl-4-methyl-cyclopentyl)-acetic
acid;
[1S-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-ethyl-4-methyl-cyclopentyl)-acetic
5 acid;
[1R-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-isopropyl-4-methyl-cyclopentyl)-
acetic acid;
[1S-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-isopropyl-4-methyl-cyclopentyl)-
acetic acid;
10 [1R-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-ethyl-4-isopropyl-cyclopentyl)-acetic
acid;
[1S-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-ethyl-4-isopropyl-cyclopentyl)-acetic
acid;
[1R-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-tert-butyl-4-methyl-cyclopentyl)-
15 acetic acid;
[1S-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-tert-butyl-4-methyl-cyclopentyl)-
acetic acid;
[1R-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-tert-butyl-4-ethyl-cyclopentyl)-acetic
acid;
20 [1S-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-tert-butyl-4-ethyl-cyclopentyl)-acetic
acid;
[1R-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-tert-butyl-4-isopropyl-cyclopentyl)-
acetic acid;
[1S-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-tert-butyl-4-isopropyl-cyclopentyl)-
25 acetic acid;
(1 α ,3 β ,4 β)-(1-Aminomethyl-3,4-di-tert-butyl-cyclopentyl)-acetic acid;
[1R-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-methyl-4-phenyl-cyclopentyl)-acetic
acid;
[1S-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-methyl-4-phenyl-cyclopentyl)-acetic
30 acid;

-7-

[1R-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-benzyl-4-methyl-cyclopentyl)-acetic acid;

[1S-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-benzyl-4-methyl-cyclopentyl)-acetic acid;

- 5 (1R-trans)-(1-Aminomethyl-3-methyl-cyclopentyl)-acetic acid;
(1R-trans)-(1-Aminomethyl-3-ethyl-cyclopentyl)-acetic acid;
(1R-trans)-(1-Aminomethyl-3-isopropyl-cyclopentyl)-acetic acid;
(1R-trans)-(1-Aminomethyl-3-tert-butyl-cyclopentyl)-acetic acid;
(1R-trans)-(1-Aminomethyl-3-phenyl-cyclopentyl)-acetic acid;
- 10 (1R-trans)-(1-Aminomethyl-3-benzyl-cyclopentyl)-acetic acid;
(1S-trans)-(1-Aminomethyl-3-methyl-cyclopentyl)-acetic acid;
(1S-trans)-(1-Aminomethyl-3-ethyl-cyclopentyl)-acetic acid;
(1S-trans)-(1-Aminomethyl-3-isopropyl-cyclopentyl)-acetic acid;
(1S-trans)-(1-Aminomethyl-3-tert-butyl-cyclopentyl)-acetic acid;
- 15 (1S-trans)-(1-Aminomethyl-3-phenyl-cyclopentyl)-acetic acid;
(1S-trans)-(1-Aminomethyl-3-benzyl-cyclopentyl)-acetic acid;
(R)-(1-Aminomethyl-3,3-dimethyl-cyclopentyl)-acetic acid;
(R)-(1-Aminomethyl-3,3-diethyl-cyclopentyl)-acetic acid;
cis-(1-Aminomethyl-3-methyl-cyclobutyl)-acetic acid;
- 20 cis-(1-Aminomethyl-3-ethyl-cyclobutyl)-acetic acid;
cis-(1-Aminomethyl-3-isopropyl-cyclobutyl)-acetic acid;
cis-(1-Aminomethyl-3-tert-butyl-cyclobutyl)-acetic acid;
cis-(1-Aminomethyl-3-phenyl-cyclobutyl)-acetic acid;
cis-(1-Aminomethyl-3-benzyl-cyclobutyl)-acetic acid;
- 25 trans-(1-Aminomethyl-3-methyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-ethyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-isopropyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-tert-butyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-phenyl-cyclobutyl)-acetic acid;
- 30 trans-(1-Aminomethyl-3-benzyl-cyclobutyl)-acetic acid;
cis-(1-Aminomethyl-3-ethyl-3-methyl-cyclobutyl)-acetic acid;

-8-

- cis-(1-Aminomethyl-3-isopropyl-3-methyl-cyclobutyl)-acetic acid;
cis-(1-Aminomethyl-3-tert-butyl-3-methyl-cyclobutyl)-acetic acid;
cis-(1-Aminomethyl-3-methyl-3-phenyl-cyclobutyl)-acetic acid;
cis-(1-Aminomethyl-3-benzyl-3-methyl-cyclobutyl)-acetic acid;
5 trans-(1-Aminomethyl-3-ethyl-3-methyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-isopropyl-3-methyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-tert-butyl-3-methyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-methyl-3-phenyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-benzyl-3-methyl-cyclobutyl)-acetic acid;
10 cis-(1-Aminomethyl-3-ethyl-3-isopropyl-cyclobutyl)-acetic acid;
cis-(1-Aminomethyl-3-tert-butyl-3-ethyl-cyclobutyl)-acetic acid;
cis-(1-Aminomethyl-3-ethyl-3-phenyl-cyclobutyl)-acetic acid;
cis-(1-Aminomethyl-3-benzyl-3-ethyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-ethyl-3-isopropyl-cyclobutyl)-acetic acid;
15 trans-(1-Aminomethyl-3-tert-butyl-3-ethyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-ethyl-3-phenyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-benzyl-3-ethyl-cyclobutyl)-acetic acid;
cis-(1-Aminomethyl-3-tert-butyl-3-isopropyl-cyclobutyl)-acetic acid;
cis-(1-Aminomethyl-3-isopropyl-3-phenyl-cyclobutyl)-acetic acid;
20 trans-(1-Aminomethyl-3-benzyl-3-isopropyl-cyclobutyl)-acetic acid;
cis-(1-Aminomethyl-3-tert-butyl-3-phenyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-benzyl-3-tert-butyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-tert-butyl-3-isopropyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-isopropyl-3-phenyl-cyclobutyl)-acetic acid;
25 cis-(1-Aminomethyl-3-benzyl-3-isopropyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-tert-butyl-3-phenyl-cyclobutyl)-acetic acid;
cis-(1-Aminomethyl-3-benzyl-3-tert-butyl-cyclobutyl)-acetic acid;
(1-Aminomethyl-3,3-dimethyl-cyclobutyl)-acetic acid;
(1-Aminomethyl-3,3-diethyl-cyclobutyl)-acetic acid;
30 (1-Aminomethyl-3,3-diisopropyl-cyclobutyl)-acetic acid;
(1-Aminomethyl-3,3-di-tert-butyl-cyclobutyl)-acetic acid;
(1-Aminomethyl-3,3-diphenyl-cyclobutyl)-acetic acid;

-9-

- (1-Aminomethyl-3,3-dibenzyl-cyclobutyl)-acetic acid;
 (1-Aminomethyl-2,2,4,4-tetramethyl-cyclobutyl)-acetic acid;
 (1-Aminomethyl-2,2,3,3,4,4-hexamethyl-cyclobutyl)-acetic acid;
 (R)-(1-Aminomethyl-2,2-dimethyl-cyclobutyl)-acetic acid;
 5 (S)-(1-Aminomethyl-2,2-dimethyl-cyclobutyl)-acetic acid;
 (1R-cis)-(1-Aminomethyl-2-methyl-cyclobutyl)-acetic acid;
 [1R-(1 α ,2 α ,3 α)]-(1-Aminomethyl-2,3-dimethyl-cyclobutyl)-acetic acid;
 (1 α ,2 α ,4 α)-(1-Aminomethyl-2,4-dimethyl-cyclobutyl)-acetic acid;
 [1R-(1 α ,2 α ,3 β)]-(1-Aminomethyl-2,3-dimethyl-cyclobutyl)-acetic acid;
 10 (1 α ,2 α ,4 β)-(1-Aminomethyl-2,4-dimethyl-cyclobutyl)-acetic acid;
 (1S-trans)-(1-Aminomethyl-2-methyl-cyclobutyl)-acetic acid;
 [1S-(1 α ,2 β ,3 β)]-(1-Aminomethyl-2,3-dimethyl-cyclobutyl)-acetic acid;
 (1 α ,2 β ,4 β)-(1-Aminomethyl-2,4-dimethyl-cyclobutyl)-acetic acid;
 [1S-(1 α ,2 β ,3 α)]-(1-Aminomethyl-2,3-dimethyl-cyclobutyl)-acetic acid;
 15 (1 α ,2 β ,4 α)-(1-Aminomethyl-2,4-dimethyl-cyclobutyl)-acetic acid;
 (1R-trans)-(1-Aminomethyl-2-methyl-cyclobutyl)-acetic acid;
 [1R-(1 α ,2 β ,3 β)]-(1-Aminomethyl-2,3-dimethyl-cyclobutyl)-acetic acid;
 [1R-(1 α ,2 β ,4 β)]-(1-Aminomethyl-2-ethyl-4-methyl-cyclobutyl)-acetic
 acid;
 20 [1R-(1 α ,2 β ,3 α)]-(1-Aminomethyl-2,3-dimethyl-cyclobutyl)-acetic acid;
 (1 α ,2 β ,4 α)-(1-Aminomethyl-2,4-dimethyl-cyclobutyl)-acetic acid;
 (1S-cis)-(1-Aminomethyl-2-methyl-cyclobutyl)-acetic acid;
 [1S-(1 α ,2 α ,3 α)]-(1-Aminomethyl-2,3-dimethyl-cyclobutyl)-acetic acid;
 [1S-(1 α ,2 α ,3 α)]-(1-Aminomethyl-2,4-dimethyl-cyclobutyl)-acetic acid;
 25 [1S-(1 α ,2 β ,3 α)]-(1-Aminomethyl-2,3-dimethyl-cyclobutyl)-acetic acid;
 (1 α ,2 α ,4 β)-(1-Aminomethyl-2,4-dimethyl-cyclobutyl)-acetic acid;
 (3S, 4S)-(1-Aminomethyl-3,4-dimethyl-cyclopentyl)-acetic acid;
 (3R, 4R)-(1-Aminomethyl-3,4-dimethyl-cyclopentyl)-acetic acid;
 (3R, 4R)-(1-Aminomethyl-3,4-diethyl-cyclopentyl)-acetic acid;
 30 (3S, 4S)-(1-Aminomethyl-3,4-diisopropyl-cyclopentyl)-acetic acid;

-10-

- (3R, 4R)-(1-Aminomethyl-3,4-diisopropyl-cyclopentyl)-acetic acid;
(3S, 4S)-(1-Aminomethyl-3,4-di-tert-butyl-cyclopentyl)-acetic acid;
(3R, 4R)-(1-Aminomethyl-3,4-di-tert-butyl-cyclopentyl)-acetic acid;
(3S, 4S)-(1-Aminomethyl-3,4-diphenyl-cyclopentyl)-acetic acid;
5 (3R, 4R)-(1-Aminomethyl-3,4-diphenyl-cyclopentyl)-acetic acid;
(3S, 4S)-(1-Aminomethyl-3,4-dibenzyl-cyclopentyl)-acetic acid;
(3R, 4R)-(1-Aminomethyl-3,4-dibenzyl-cyclopentyl)-acetic acid;
[1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-methyl-4-ethyl-cyclopentyl)-acetic
acid;
10 [1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-methyl-4-ethyl-cyclopentyl)-acetic
acid;
[1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-methyl-4-ethyl-cyclopentyl)-acetic
acid;
[1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-methyl-4-ethyl-cyclopentyl)-acetic
15 acid;
[1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-methyl-4-isopropyl-cyclopentyl)-
acetic acid;
[1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-methyl-4-isopropyl-cyclopentyl)-
acetic acid;
20 [1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-methyl-4-isopropyl-cyclopentyl)-
acetic acid;
[1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-methyl-4-isopropyl-cyclopentyl)-
acetic acid;
[1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-methyl-4-tert-butyl-cyclopentyl)-
25 acetic acid;
[1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-methyl-4-tert-butyl-cyclopentyl)-
acetic acid;
[1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-methyl-4-tert-butyl-cyclopentyl)-
acetic acid;

-11-

- [1S-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-methyl-4-tert-butyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-methyl-4-phenyl-cyclopentyl)-acetic acid;
- 5 [1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-methyl-4-phenyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-methyl-4-phenyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-methyl-4-phenyl-cyclopentyl)-acetic acid;
- 10 [1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-benzyl-4-methyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-benzyl-4-methyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-benzyl-4-methyl-cyclopentyl)-acetic acid;
- 15 [1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-benzyl-4-methyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-benzyl-4-methyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-ethyl-4-isopropyl-cyclopentyl)-acetic acid;
- 20 [1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-ethyl-4-isopropyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-ethyl-4-isopropyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-ethyl-4-isopropyl-cyclopentyl)-acetic acid;
- 25 [1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-ethyl-4-isopropyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-tert-butyl-4-ethyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-tert-butyl-4-ethyl-cyclopentyl)-acetic acid;
- 30 [1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-tert-butyl-4-ethyl-cyclopentyl)-acetic acid;

-12-

- [1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-tert-butyl-4-ethyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-tert-butyl-4-ethyl-cyclopentyl)-acetic acid;
- 5 [1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-ethyl-4-phenyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-ethyl-4-phenyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-ethyl-4-phenyl-cyclopentyl)-acetic acid;
- 10 [1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-ethyl-4-phenyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-benzyl-4-ethyl-cyclopentyl)-acetic acid;
- 15 [1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-benzyl-4-ethyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-benzyl-4-ethyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-benzyl-4-ethyl-cyclopentyl)-acetic acid;
- 20 [1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-tert-butyl-4-isopropyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-tert-butyl-4-isopropyl-cyclopentyl)-acetic acid;
- 25 [1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-tert-butyl-4-isopropyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-tert-butyl-4-isopropyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-isopropyl-4-phenyl-cyclopentyl)-acetic acid;
- 30 acetic acid;

-13-

- [1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-isopropyl-4-phenyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-isopropyl-4-phenyl-cyclopentyl)-acetic acid;
- 5 [1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-isopropyl-4-phenyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-benzyl-4-isopropyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-benzyl-4-isopropyl-cyclopentyl)-acetic acid;
- 10 [1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-benzyl-4-isopropyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-benzyl-4-isopropyl-cyclopentyl)-acetic acid;
- 15 [1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-tert-butyl-4-phenyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-tert-butyl-4-phenyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-tert-butyl-4-phenyl-cyclopentyl)-acetic acid;
- 20 [1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-tert-butyl-4-phenyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-benzyl-4-tert-butyl-cyclopentyl)-acetic acid;
- 25 [1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-benzyl-4-tert-butyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-benzyl-4-tert-butyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-benzyl-4-tert-butyl-cyclopentyl)-acetic acid;
- 30

-14-

- [1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-benzyl-4-phenyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-benzyl-4-phenyl-cyclopentyl)-acetic acid;
- 5 [1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-benzyl-4-phenyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-benzyl-4-phenyl-cyclopentyl)-acetic acid;
- (1R-cis)-(1-Aminomethyl-2-methyl-cyclopentyl)-acetic acid;
- 10 (1S-cis)-(1-Aminomethyl-2-methyl-cyclopentyl)-acetic acid;
- (1R-trans)-(1-Aminomethyl-2-methyl-cyclopentyl)-acetic acid;
- (1S-trans)-(1-Aminomethyl-2-methyl-cyclopentyl)-acetic acid;
- (R)-(1-Aminomethyl-2,2-dimethyl-cyclopentyl)-acetic acid;
- (S)-(1-Aminomethyl-2,2-dimethyl-cyclopentyl)-acetic acid;
- 15 (1-Aminomethyl-2,2,5,5-tetramethyl-cyclopentyl)-acetic acid;
- (1 α ,2 β ,5 β)-(1-Aminomethyl-2,5-dimethyl-cyclopentyl)-acetic acid;
- (2R, 5R)-(1-Aminomethyl-2,5-dimethyl-cyclopentyl)-acetic acid;
- (2S, 5S)-(1-Aminomethyl-2,5-dimethyl-cyclopentyl)-acetic acid;
- (1 α ,2 α ,5 α)-(1-Aminomethyl-2,5-dimethyl-cyclopentyl)-acetic acid;
- 20 [1R-(1 α ,2 α ,3 α)]-(1-Aminomethyl-2,3-dimethyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,2 β ,3 α)]-(1-Aminomethyl-2,3-dimethyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,2 α ,3 β)]-(1-Aminomethyl-2,3-dimethyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,2 β ,3 β)]-(1-Aminomethyl-2,3-dimethyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,2 α ,3 α)]-(1-Aminomethyl-2,3-dimethyl-cyclopentyl)-acetic acid;
- 25 [1S-(1 α ,2 β ,3 α)]-(1-Aminomethyl-2,3-dimethyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,2 α ,3 β)]-(1-Aminomethyl-2,3-dimethyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,2 β ,3 β)]-(1-Aminomethyl-2,3-dimethyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,2 α ,4 α)]-(1-Aminomethyl-2,4-dimethyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,2 α ,4 α)]-(1-Aminomethyl-2,4-dimethyl-cyclopentyl)-acetic acid;
- 30 [1R-(1 α ,2 α ,4 β)]-(1-Aminomethyl-2,4-dimethyl-cyclopentyl)-acetic acid;

-15-

[1S-(1 α ,2 α ,4 β)]-(1-Aminomethyl-2,4-dimethyl-cyclopentyl)-acetic acid;

[1R-(1 α ,2 β ,4 α)]-(1-Aminomethyl-2,4-dimethyl-cyclopentyl)-acetic acid;

[1S-(1 α ,2 β ,4 α)]-(1-Aminomethyl-2,4-dimethyl-cyclopentyl)-acetic acid;

[1R-(1 α ,2 β ,4 β)]-(1-Aminomethyl-2,4-dimethyl-cyclopentyl)-acetic acid;

5 and

[1S-(1 α ,2 β ,4 β)]-(1-Aminomethyl-2,4-dimethyl-cyclopentyl)-acetic acid.

Especially useful as an agent for insomnia and related disorders is

(3S, 4S)-(1-Aminomethyl-3,4-dimethyl-cyclopentyl)-acetic acid.

Also, especially useful as an agent for insomnia and related disorders is

10 (3S, 5R)-3-Aminomethyl-5-methyl-octanoic acid.

The term "lower alkyl" is a straight or branched group of from 1 to 4 carbons.

The term "alkyl" is a straight or branched group of from 1 to 6 carbon atoms including but not limited to methyl, ethyl, propyl, n-propyl, isopropyl, butyl, 2-butyl, tert-butyl, pentyl, except as where otherwise stated.

15

The benzyl and phenyl groups may be unsubstituted or substituted by from 1 to 3 substituents selected from hydroxy, carboxy, carboalkoxy, halogen, CF₃, nitro, alkyl, and alkoxy. Preferred are halogens.

Since amino acids are amphoteric, pharmacologically compatible salts when R is hydrogen can be salts of appropriate inorganic or organic acids, for example, hydrochloric, sulphuric, phosphoric, acetic, oxalic, lactic, citric, malic, salicylic, malonic, maleic, succinic, methanesulfonic acid, and ascorbic. Starting from corresponding hydroxides or carbonates, salts with alkali metals or alkaline earth metals, for example, sodium, potassium, magnesium, or calcium are formed. Salts with quaternary ammonium ions can also be prepared with, for example, the tetramethyl-ammonium ion. The carboxyl group of the amino acids can be esterified by known means.

25

Certain of the compounds of the present invention can exist in unsolvated forms as well as solvated forms, including hydrated forms. In general, the solvated forms, including hydrated forms, are equivalent to unsolvated forms and are intended to be encompassed within the scope of the present invention.

30

-16-

Certain of the compounds of the present invention possess one or more chiral centers and each center may exist in the R(D) or S(L) configuration. The present invention includes all enantiomeric and epimeric forms as well as the appropriate mixtures thereof.

5

METHODS AND MATERIALS

Methods

Male Sprague-Dawley rats weighing 250 to 360 g were used for the experiment. The rats were kept on a 12:12-hr light/dark cycle (lights on at 0900) at $23 \pm 1^\circ\text{C}$ ambient temperature. They had free access water and food during the experiment. Stainless steel jewelry screws for EEG recording were placed over the frontal and parietal cortices. An EMG electrode was implanted in the dorsal neck muscles.

Recording and analyses. After the recovery period (at least 1 week), rats were moved to sleep recording chambers. The rats were allowed relatively unrestricted movement inside the recording cages. A flexible tether connected the electrodes led to an electronic swivel. The leads from the swivel were routed to Grass Model No. 7D polygraphs in an adjacent room for recording EEG and EMG activity. The vigilance states of wakefulness, nonrapid-eye-movement sleep (NREMS), and rapid-eye-movement sleep (REMS) were determined offline in 10 second epochs. The amount of time spent in each vigilance state was calculated every hour and totaled for the 12-hour period postinjection. Drug or vehicle were administered PO just prior to the onset of the light phase of the rat's light/dark cycle. Each rat received a vehicle and one drug dose.

Results

(3S, 4S)-1-Aminomethyl-3,4-dimethyl-cyclopentyl)-acetic acid (Compound 1), 3, 10, and 30 mg/kg PO and (3S, 5R)-3-Aminomethyl-5-methyl-octanoic acid (Compound 2), 1, 3, and 20 mg/kg PO significantly increased NREMS in rats during the 12-hour period following injection (Table 1 and

-17-

Table 2, respectively). This increase in NREMS sleep in rats is the typical profile demonstrated by agents that are used as sleep-inducing agents in humans. (Meltzer L.T. and Serpa K.A., Assessment of hypnotic effects in the rat: Influence of the sleep-awake cycle, *Drug Development Research* 1988;14:151-159;

5 Depoortere H. et al., Hypnotics: Clinical value of pharmaco-EEG methods, *Neuropsychobiology* 1986;16:157-162).

TABLE 1. Compound 1 Increases NREMS Sleep in Rats (N = 8/Treatment)

Total Minutes (Mean \pm SEM) NREMS for 12-Hour Postinjection		
Compound 1 (mg/kg PO)	Vehicle	Drug
3	370 \pm 11	401 \pm 8*
10	383 \pm 8	409 \pm 12*
30	371 \pm 14	424 \pm 6*

* P \leq 0.05 vs Vehicle (paired t-test)

TABLE 2. Compound 2 Increases NREMS Sleep in Rats (N = 8/Treatment)

Total Minutes (Mean \pm SEM) NREMS for 12-Hour Postinjection		
Compound 2 (mg/kg PO)	Vehicle	Drug
1	409 \pm 8	432 \pm 8*
3	406 \pm 9	451 \pm 9*
20	412 \pm 6	481 \pm 6*

* P \leq 0.05 vs Vehicle (paired t-test)

The compounds of the present invention can be prepared and administered in a wide variety of oral and parenteral dosage forms. Thus, the compounds of the present invention can be administered by injection, that is, intravenously,

10 intramuscularly, intracutaneously, subcutaneously, intraduodenally, or intraperitoneally. Also, the compounds of the present invention can be administered by inhalation, for example, intranasally. Additionally, the compounds of the present invention can be administered transdermally. It will be

-18-

obvious to those skilled in the art that the following dosage forms may comprise as the active component, either a compound of formula 1 or 1A or a corresponding pharmaceutically acceptable salt of a compound of formula 1 or 1A.

5 For preparing pharmaceutical compositions from the compounds of the present invention, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, pills, capsules, cachets, suppositories, and dispersible granules. A solid carrier can be one or more substances which may also act as diluents, flavoring agents, binders, preservatives, tablet disintegrating agents, or an encapsulating material.

10 In powders, the carrier is a finely divided solid which is in a mixture with the finely divided active component.

In tablets, the active component is mixed with the carrier having the necessary binding properties in suitable proportions and compacted in the shape and size desired.

15 The powders and tablets preferably contain from five or ten to about seventy percent of the active compound. Suitable carriers are magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter, and the like. The term "preparation" is intended to include the
20 formulation of the active compound with encapsulating material as a carrier providing a capsule in which the active component, with or without other carriers, is surrounded by a carrier, which is thus in association with it. Similarly, cachets and lozenges are included. Tablets, powders, capsules, pills, cachets, and lozenges can be used as solid dosage forms suitable for oral administration.

25 For preparing suppositories, a low melting wax, such as a mixture of fatty acid glycerides or cocoa butter, is first melted, and the active component is dispersed homogeneously therein, as by stirring. The molten homogenous mixture is then poured into convenient sized molds, allowed to cool, and thereby to solidify.

30 Liquid form preparations include solutions, suspensions, and emulsions, for example, water or water propylene glycol solutions. For parenteral injection

-19-

liquid preparations can be formulated in solution in aqueous polyethylene glycol solution.

Aqueous solutions suitable for oral use can be prepared by dissolving the active component in water and adding suitable colorants, flavors, stabilizing and thickening agents as desired.

Aqueous suspensions suitable for oral use can be made by dispersing the finely divided active component in water with viscous material, such as natural or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, and other well-known suspending agents.

Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations for oral administration. Such liquid forms include solutions, suspensions, and emulsions. These preparations may contain, in addition to the active component, colorants, flavors, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like.

The pharmaceutical preparation is preferably in unit dosage form. In such form the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as packeted tablets, capsules, and powders in vials or ampoules. Also, the unit dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form.

The quantity of active component in a unit dose preparation may be varied or adjusted from 0.1 mg to 1 g according to the particular application and the potency of the active component. In medical use the drug may be administered three times daily as, for example, capsules of 100 or 300 mg. The composition can, if desired, also contain other compatible therapeutic agents.

In therapeutic use, the compounds utilized in the pharmaceutical method of this invention are administered at the initial dosage of about 0.01 mg to about 100 mg/kg daily. A daily dose range of about 0.01 mg to about 100 mg/kg is preferred. The dosages, however, may be varied depending upon the requirements of the patient, the severity of the condition being treated, and the compound being


-20-

employed. Determination of the proper dosage for a particular situation is within the skill of the art. Generally, treatment is initiated with smaller dosages which are less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the optimum effect under the circumstances is reached.

- 5 For convenience, the total daily dosage may be divided and administered in portions during the day, if desired.

CLAIMS

1. A method of treating sleep disorders in a mammal in need of said treatment comprising administering a therapeutically effective amount of a compound of formula



or a pharmaceutically acceptable salt thereof wherein:

R is hydrogen or a lower alkyl;

R¹ to R¹⁴ are each independently selected from hydrogen, straight or branched alkyl of from 1 to 6 carbons, phenyl, benzyl, fluorine, chlorine, bromine, hydroxy, hydroxymethyl, amino, aminomethyl, trifluoromethyl, -CO₂H, -CO₂R¹⁵, -CH₂CO₂H, -CH₂CO₂R¹⁵, -OR¹⁵ wherein R¹⁵ is a straight or branched alkyl of from 1 to 6 carbons, phenyl, or benzyl, and R¹ to R⁸ are not simultaneously hydrogen.

- 15 2. A method according to Claim 1 wherein R¹ to R¹⁴ are selected from
hydrogen, methyl, ethyl, propyl, isopropyl, butyl straight or branched,
phenyl, or benzyl.
3. A method according to Claim 1 wherein R¹ to R¹⁴ are selected from
hydrogen, methyl, ethyl, or benzyl.
- 20 4. A method according to Claim 1 wherein the compound administered is
selected from:

-22-

- (±)-(trans)-(1-Aminomethyl-3,4-dimethyl-cyclopentyl)-acetic acid hydrochloride;
(1-Aminomethyl-cyclobutyl)-acetic acid hydrochloride;
(cis/trans)-(3R)-(1-Aminomethyl-3-methyl-cyclopentyl)-acetic acid hydrochloride;
5 (cis/trans)-(1-Aminomethyl-3-benzyl-cyclobutyl)-acetic acid hydrochloride;
(±)-(1-Aminomethyl-3,3-dimethyl-cyclopentyl)-acetic acid hydrochloride; and
10 (cis)-(3R)-(1-Aminomethyl-3-methyl-cyclopentyl)-acetic acid hydrochloride.
5. A method according to Claim 1 wherein the compound administered is selected from:
- 15 (1 α ,3 α ,4 α)-(1-Aminomethyl-3,4-dimethyl-cyclopentyl)-acetic acid;
(1 α ,3 α ,4 α)-(1-Aminomethyl-3,4-diethyl-cyclopentyl)-acetic acid;
(1 α ,3 α ,4 α)-(1-Aminomethyl-3,4-diisopropyl-cyclopentyl)-acetic acid;
[1S-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-ethyl-4-methyl-cyclopentyl)-acetic acid;
20 [1R-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-ethyl-4-methyl-cyclopentyl)-acetic acid;
[1S-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-isopropyl-4-methyl-cyclopentyl)-acetic acid;
[1R-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-isopropyl-4-methyl-cyclopentyl)-acetic acid;
25 [1S-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-ethyl-4-isopropyl-cyclopentyl)-acetic acid;
[1R-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-ethyl-4-isopropyl-cyclopentyl)-acetic acid;

-23-

- [1S-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-tert-butyl-4-methyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-tert-butyl-4-methyl-cyclopentyl)-acetic acid;
- 5 [1S-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-tert-butyl-4-ethyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-tert-butyl-4-ethyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-tert-butyl-4-isopropyl-cyclopentyl)-acetic acid;
- 10 [1R-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-tert-butyl-4-isopropyl-cyclopentyl)-acetic acid;
- (1 α ,3 α ,4 α)-(1-Aminomethyl-3,4-di-tert-butyl-cyclopentyl)-acetic acid;
- 15 [1S-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-methyl-4-phenyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-methyl-4-phenyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-benzyl-4-methyl-cyclopentyl)-acetic acid;
- 20 [1R-(1 α ,3 α ,4 α)]-(1-Aminomethyl-3-benzyl-4-methyl-cyclopentyl)-acetic acid;
- (1S-cis)-(1-Aminomethyl-3-methyl-cyclopentyl)-acetic acid;
- (1S-cis)-(1-Aminomethyl-3-ethyl-cyclopentyl)-acetic acid;
- 25 (1S-cis)-(1-Aminomethyl-3-isopropyl-cyclopentyl)-acetic acid;
- (1S-cis)-(1-Aminomethyl-3-tert-butyl-cyclopentyl)-acetic acid;
- (1S-cis)-(1-Aminomethyl-3-phenyl-cyclopentyl)-acetic acid;
- (1S-cis)-(1-Aminomethyl-3-benzyl-cyclopentyl)-acetic acid;
- (1R-cis)-(1-Aminomethyl-3-methyl-cyclopentyl)-acetic acid;
- 30 (1R-cis)-(1-Aminomethyl-3-ethyl-cyclopentyl)-acetic acid;
- (1R-cis)-(1-Aminomethyl-3-isopropyl-cyclopentyl)-acetic acid;

-24-

- (1R-cis)-(1-Aminomethyl-3-tert-butyl-cyclopentyl)-acetic acid;
(1R-cis)-(1-Aminomethyl-3-phenyl-cyclopentyl)-acetic acid;
(1R-cis)-(1-Aminomethyl-3-benzyl-cyclopentyl)-acetic acid;
(S)-(1-Aminomethyl-3,3-dimethyl-cyclopentyl)-acetic acid;
5 (S)-(1-Aminomethyl-3,3-diethyl-cyclopentyl)-acetic acid;
(1-Aminomethyl-3,3,4,4-tetramethyl-cyclopentyl)-acetic acid;
(1-Aminomethyl-3,3,4,4-tetraethyl-cyclopentyl)-acetic acid;
(1 α ,3 β ,4 β)-(1-Aminomethyl-3,4-dimethyl-cyclopentyl)-acetic acid;
(1 α ,3 β ,4 β)-(1-Aminomethyl-3,4-diethyl-cyclopentyl)-acetic acid;
10 (1 α ,3 β ,4 β)-(1-Aminomethyl-3,4-diisopropyl-cyclopentyl)-acetic
acid;
[1R-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-ethyl-4-methyl-cyclopentyl)-
acetic acid;
[1S-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-ethyl-4-methyl-cyclopentyl)-
15 acetic acid;
[1R-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-isopropyl-4-methyl-
cyclopentyl)-acetic acid;
[1S-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-isopropyl-4-methyl-
cyclopentyl)-acetic acid;
20 [1R-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-ethyl-4-isopropyl-
cyclopentyl)-acetic acid;
[1S-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-ethyl-4-isopropyl-cyclopentyl)-
acetic acid;
[1R-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-tert-butyl-4-methyl-
25 cyclopentyl)-acetic acid;
[1S-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-tert-butyl-4-methyl-
cyclopentyl)-acetic acid;
[1R-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-tert-butyl-4-ethyl-
cyclopentyl)-acetic acid;

-25-

- [1S-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-tert-butyl-4-ethyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-tert-butyl-4-isopropyl-cyclopentyl)-acetic acid;
- 5 [1S-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-tert-butyl-4-isopropyl-cyclopentyl)-acetic acid;
- (1 α ,3 β ,4 β)-(1-Aminomethyl-3,4-di-tert-butyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-methyl-4-phenyl-cyclopentyl)-acetic acid;
- 10 [1S-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-methyl-4-phenyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-benzyl-4-methyl-cyclopentyl)-acetic acid;
- 15 [1S-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-benzyl-4-methyl-cyclopentyl)-acetic acid;
- (1R-trans)-(1-Aminomethyl-3-methyl-cyclopentyl)-acetic acid;
- (1R-trans)-(1-Aminomethyl-3-ethyl-cyclopentyl)-acetic acid;
- (1R-trans)-(1-Aminomethyl-3-isopropyl-cyclopentyl)-acetic acid;
- 20 (1R-trans)-(1-Aminomethyl-3-tert-butyl-cyclopentyl)-acetic acid;
- (1R-trans)-(1-Aminomethyl-3-phenyl-cyclopentyl)-acetic acid;
- (1R-trans)-(1-Aminomethyl-3-benzyl-cyclopentyl)-acetic acid;
- (1S-trans)-(1-Aminomethyl-3-methyl-cyclopentyl)-acetic acid;
- (1S-trans)-(1-Aminomethyl-3-ethyl-cyclopentyl)-acetic acid;
- 25 (1S-trans)-(1-Aminomethyl-3-isopropyl-cyclopentyl)-acetic acid;
- (1S-trans)-(1-Aminomethyl-3-tert-butyl-cyclopentyl)-acetic acid;
- (1S-trans)-(1-Aminomethyl-3-phenyl-cyclopentyl)-acetic acid;
- (1S-trans)-(1-Aminomethyl-3-benzyl-cyclopentyl)-acetic acid;
- (R)-(1-Aminomethyl-3,3-dimethyl-cyclopentyl)-acetic acid;
- 30 (R)-(1-Aminomethyl-3,3-diethyl-cyclopentyl)-acetic acid;
- cis-(1-Aminomethyl-3-methyl-cyclobutyl)-acetic acid;

-26-

- 5 cis-(1-Aminomethyl-3-ethyl-cyclobutyl)-acetic acid;
cis-(1-Aminomethyl-3-isopropyl-cyclobutyl)-acetic acid;
cis-(1-Aminomethyl-3-tert-butyl-cyclobutyl)-acetic acid;
cis-(1-Aminomethyl-3-phenyl-cyclobutyl)-acetic acid;
cis-(1-Aminomethyl-3-benzyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-methyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-ethyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-isopropyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-tert-butyl-cyclobutyl)-acetic acid;
10 trans-(1-Aminomethyl-3-phenyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-benzyl-cyclobutyl)-acetic acid;
cis-(1-Aminomethyl-3-ethyl-3-methyl-cyclobutyl)-acetic acid;
cis-(1-Aminomethyl-3-isopropyl-3-methyl-cyclobutyl)-acetic acid;
cis-(1-Aminomethyl-3-tert-butyl-3-methyl-cyclobutyl)-acetic acid;
15 cis-(1-Aminomethyl-3-methyl-3-phenyl-cyclobutyl)-acetic acid;
cis-(1-Aminomethyl-3-benzyl-3-methyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-ethyl-3-methyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-isopropyl-3-methyl-cyclobutyl)-acetic
acid;
20 trans-(1-Aminomethyl-3-tert-butyl-3-methyl-cyclobutyl)-acetic
acid;
trans-(1-Aminomethyl-3-methyl-3-phenyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-benzyl-3-methyl-cyclobutyl)-acetic acid;
cis-(1-Aminomethyl-3-ethyl-3-isopropyl-cyclobutyl)-acetic acid;
25 cis-(1-Aminomethyl-3-tert-butyl-3-ethyl-cyclobutyl)-acetic acid;
cis-(1-Aminomethyl-3-ethyl-3-phenyl-cyclobutyl)-acetic acid;
cis-(1-Aminomethyl-3-benzyl-3-ethyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-ethyl-3-isopropyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-tert-butyl-3-ethyl-cyclobutyl)-acetic acid;
30 trans-(1-Aminomethyl-3-ethyl-3-phenyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-benzyl-3-ethyl-cyclobutyl)-acetic acid;

-27-

- acid;
cis-(1-Aminomethyl-3-tert-butyl-3-isopropyl-cyclobutyl)-acetic acid;
cis-(1-Aminomethyl-3-isopropyl-3-phenyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-benzyl-3-isopropyl-cyclobutyl)-acetic
5 acid;
cis-(1-Aminomethyl-3-tert-butyl-3-phenyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-benzyl-3-tert-butyl-cyclobutyl)-acetic
acid;
trans-(1-Aminomethyl-3-tert-butyl-3-isopropyl-cyclobutyl)-acetic
10 acid;
trans-(1-Aminomethyl-3-isopropyl-3-phenyl-cyclobutyl)-acetic
acid;
cis-(1-Aminomethyl-3-benzyl-3-isopropyl-cyclobutyl)-acetic acid;
trans-(1-Aminomethyl-3-tert-butyl-3-phenyl-cyclobutyl)-acetic
15 acid;
cis-(1-Aminomethyl-3-benzyl-3-tert-butyl-cyclobutyl)-acetic acid;
(1-Aminomethyl-3,3-dimethyl-cyclobutyl)-acetic acid;
(1-Aminomethyl-3,3-diethyl-cyclobutyl)-acetic acid;
(1-Aminomethyl-3,3-diisopropyl-cyclobutyl)-acetic acid;
20 (1-Aminomethyl-3,3-di-tert-butyl-cyclobutyl)-acetic acid;
(1-Aminomethyl-3,3-diphenyl-cyclobutyl)-acetic acid;
(1-Aminomethyl-3,3-dibenzyl-cyclobutyl)-acetic acid;
(1-Aminomethyl-2,2,4,4-tetramethyl-cyclobutyl)-acetic acid;
(1-Aminomethyl-2,2,3,3,4,4-hexamethyl-cyclobutyl)-acetic acid;
25 (R)-(1-Aminomethyl-2,2-dimethyl-cyclobutyl)-acetic acid;
(S)-(1-Aminomethyl-2,2-dimethyl-cyclobutyl)-acetic acid;
(1R-cis)-(1-Aminomethyl-2-methyl-cyclobutyl)-acetic acid;
[1R-(1 α ,2 α ,3 α)]-(1-Aminomethyl-2,3-dimethyl-cyclobutyl)-acetic
acid;
30 (1 α ,2 α ,4 α)-(1-Aminomethyl-2,4-dimethyl-cyclobutyl)-acetic acid;

-28-

- [1R-(1 α ,2 α ,3 β)]-(1-Aminomethyl-2,3-dimethyl-cyclobutyl)-acetic acid;
- (1 α ,2 α ,4 β)-(1-Aminomethyl-2,4-dimethyl-cyclobutyl)-acetic acid;
- (1S-trans)-(1-Aminomethyl-2-methyl-cyclobutyl)-acetic acid;
- 5 [1S-(1 α ,2 β ,3 β)]-(1-Aminomethyl-2,3-dimethyl-cyclobutyl)-acetic acid;
- (1 α ,2 β ,4 β)-(1-Aminomethyl-2,4-dimethyl-cyclobutyl)-acetic acid;
- [1S-(1 α ,2 β ,3 α)]-(1-Aminomethyl-2,3-dimethyl-cyclobutyl)-acetic acid;
- 10 (1 α ,2 β ,4 α)-(1-Aminomethyl-2,4-dimethyl-cyclobutyl)-acetic acid;
- (1R-trans)-(1-Aminomethyl-2-methyl-cyclobutyl)-acetic acid;
- [1R-(1 α ,2 β ,3 β)]-(1-Aminomethyl-2,3-dimethyl-cyclobutyl)-acetic acid;
- [1R-(1 α ,2 β ,4 β)]-(1-Aminomethyl-2-ethyl-4-methyl-cyclobutyl)-acetic acid;
- 15 [1R-(1 α ,2 β ,3 α)]-(1-Aminomethyl-2,3-dimethyl-cyclobutyl)-acetic acid;
- (1 α ,2 β ,4 α)-(1-Aminomethyl-2,4-dimethyl-cyclobutyl)-acetic acid;
- (1S-cis)-(1-Aminomethyl-2-methyl-cyclobutyl)-acetic acid;
- 20 [1S-(1 α ,2 α ,3 α)]-(1-Aminomethyl-2,3-dimethyl-cyclobutyl)-acetic acid;
- [1S-(1 α ,2 α ,3 α)]-(1-Aminomethyl-2,4-dimethyl-cyclobutyl)-acetic acid;
- [1S-(1 α ,2 β ,3 α)]-(1-Aminomethyl-2,3-dimethyl-cyclobutyl)-acetic acid;
- 25 (1 α ,2 α ,4 β)-(1-Aminomethyl-2,4-dimethyl-cyclobutyl)-acetic acid;
- (3R, 4R)-(1-Aminomethyl-3,4-dimethyl-cyclopentyl)-acetic acid;
- (3S, 4S)-(1-Aminomethyl-3,4-diethyl-cyclopentyl)-acetic acid;
- (3R, 4R)-(1-Aminomethyl-3,4-diethyl-cyclopentyl)-acetic acid;
- 30 (3S, 4S)-(1-Aminomethyl-3,4-diisopropyl-cyclopentyl)-acetic acid;

-29-

- (3R, 4R)-(1-Aminomethyl-3,4-diisopropyl-cyclopentyl)-acetic acid;
(3S, 4S)-(1-Aminomethyl-3,4-di-tert-butyl-cyclopentyl)-acetic acid;
(3R, 4R)-(1-Aminomethyl-3,4-di-tert-butyl-cyclopentyl)-acetic
acid;
5 (3S, 4S)-(1-Aminomethyl-3,4-diphenyl-cyclopentyl)-acetic acid;
(3R, 4R)-(1-Aminomethyl-3,4-diphenyl-cyclopentyl)-acetic acid;
(3S, 4S)-(1-Aminomethyl-3,4-dibenzyl-cyclopentyl)-acetic acid;
(3R, 4R)-(1-Aminomethyl-3,4-dibenzyl-cyclopentyl)-acetic acid;
[1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-methyl-4-ethyl-cyclopentyl)-
10 acetic acid;
[1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-methyl-4-ethyl-cyclopentyl)-
acetic acid;
[1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-methyl-4-ethyl-cyclopentyl)-
acetic acid;
15 [1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-methyl-4-ethyl-cyclopentyl)-
acetic acid;
[1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-methyl-4-isopropyl-
cyclopentyl)-acetic acid;
[1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-methyl-4-isopropyl-
20 cyclopentyl)-acetic acid;
[1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-methyl-4-isopropyl-
cyclopentyl)-acetic acid;
[1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-methyl-4-isopropyl-
cyclopentyl)-acetic acid;
25 [1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-methyl-4-tert-butyl-
cyclopentyl)-acetic acid;
[1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-methyl-4-tert-butyl-
cyclopentyl)-acetic acid;
[1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-methyl-4-tert-butyl-
30 cyclopentyl)-acetic acid;

-30-

- [1S-(1 α ,3 β ,4 β)]-(1-Aminomethyl-3-methyl-4-tert-butyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-methyl-4-phenyl-cyclopentyl)-acetic acid;
- 5 [1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-methyl-4-phenyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-methyl-4-phenyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-methyl-4-phenyl-cyclopentyl)-acetic acid;
- 10 [1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-benzyl-4-methyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-benzyl-4-methyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-benzyl-4-methyl-cyclopentyl)-acetic acid;
- 15 [1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-benzyl-4-methyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-benzyl-4-methyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-ethyl-4-isopropyl-cyclopentyl)-acetic acid;
- 20 [1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-ethyl-4-isopropyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-ethyl-4-isopropyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-ethyl-4-isopropyl-cyclopentyl)-acetic acid;
- 25 [1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-tert-butyl-4-ethyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-tert-butyl-4-ethyl-cyclopentyl)-acetic acid;
- 30

-31-

- [1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-tert-butyl-4-ethyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-tert-butyl-4-ethyl-cyclopentyl)-acetic acid;
- 5 [1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-ethyl-4-phenyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-ethyl-4-phenyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-ethyl-4-phenyl-cyclopentyl)-acetic acid;
- 10 [1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-ethyl-4-phenyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-benzyl-4-ethyl-cyclopentyl)-acetic acid;
- 15 [1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-benzyl-4-ethyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-benzyl-4-ethyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-benzyl-4-ethyl-cyclopentyl)-acetic acid;
- 20 [1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-tert-butyl-4-isopropyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-tert-butyl-4-isopropyl-cyclopentyl)-acetic acid;
- 25 [1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-tert-butyl-4-isopropyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-tert-butyl-4-isopropyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-isopropyl-4-phenyl-cyclopentyl)-acetic acid;
- 30

-32-

- [1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-isopropyl-4-phenyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-isopropyl-4-phenyl-cyclopentyl)-acetic acid;
- 5 [1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-isopropyl-4-phenyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-benzyl-4-isopropyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-benzyl-4-isopropyl-cyclopentyl)-acetic acid;
- 10 [1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-benzyl-4-isopropyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-benzyl-4-isopropyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-tert-butyl-4-phenyl-cyclopentyl)-acetic acid;
- 15 [1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-tert-butyl-4-phenyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-tert-butyl-4-phenyl-cyclopentyl)-acetic acid;
- 20 [1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-tert-butyl-4-phenyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-benzyl-4-tert-butyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-benzyl-4-tert-butyl-cyclopentyl)-acetic acid;
- 25 [1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-benzyl-4-tert-butyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-benzyl-4-tert-butyl-cyclopentyl)-acetic acid;
- 30 [1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-benzyl-4-tert-butyl-cyclopentyl)-acetic acid;

-33-

- [1S-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-benzyl-4-phenyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-benzyl-4-phenyl-cyclopentyl)-acetic acid;
- 5 [1R-(1 α ,3 α ,4 β)]-(1-Aminomethyl-3-benzyl-4-phenyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,3 β ,4 α)]-(1-Aminomethyl-3-benzyl-4-phenyl-cyclopentyl)-acetic acid;
- (1R-cis)-(1-Aminomethyl-2-methyl-cyclopentyl)-acetic acid;
- 10 (1S-cis)-(1-Aminomethyl-2-methyl-cyclopentyl)-acetic acid;
- (1R-trans)-(1-Aminomethyl-2-methyl-cyclopentyl)-acetic acid;
- (1S-trans)-(1-Aminomethyl-2-methyl-cyclopentyl)-acetic acid;
- (R)-(1-Aminomethyl-2,2-dimethyl-cyclopentyl)-acetic acid;
- (S)-(1-Aminomethyl-2,2-dimethyl-cyclopentyl)-acetic acid;
- 15 (1-Aminomethyl-2,2,5,5-tetramethyl-cyclopentyl)-acetic acid;
- (1 α ,2 β ,5 β)-(1-Aminomethyl-2,5-dimethyl-cyclopentyl)-acetic acid;
- (2R, 5R)-(1-Aminomethyl-2,5-dimethyl-cyclopentyl)-acetic acid;
- (2S, 5S)-(1-Aminomethyl-2,5-dimethyl-cyclopentyl)-acetic acid;
- (1 α ,2 α ,5 α)-(1-Aminomethyl-2,5-dimethyl-cyclopentyl)-acetic acid;
- 20 [1R-(1 α ,2 α ,3 α)]-(1-Aminomethyl-2,3-dimethyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,2 β ,3 α)]-(1-Aminomethyl-2,3-dimethyl-cyclopentyl)-acetic acid;
- [1R-(1 α ,2 α ,3 β)]-(1-Aminomethyl-2,3-dimethyl-cyclopentyl)-acetic acid;
- 25 [1R-(1 α ,2 β ,3 β)]-(1-Aminomethyl-2,3-dimethyl-cyclopentyl)-acetic acid;
- [1S-(1 α ,2 α ,3 α)]-(1-Aminomethyl-2,3-dimethyl-cyclopentyl)-acetic acid;

-34-

[1S-(1 α ,2 β ,3 α)]-(1-Aminomethyl-2,3-dimethyl-cyclopentyl)-acetic acid;
[1S-(1 α ,2 α ,3 β)]-(1-Aminomethyl-2,3-dimethyl-cyclopentyl)-acetic acid;
5 [1S-(1 α ,2 β ,3 β)]-(1-Aminomethyl-2,3-dimethyl-cyclopentyl)-acetic acid;
[1R-(1 α ,2 α ,4 α)]-(1-Aminomethyl-2,4-dimethyl-cyclopentyl)-acetic acid;
[1S-(1 α ,2 α ,4 α)]-(1-Aminomethyl-2,4-dimethyl-cyclopentyl)-acetic
10 acid;
[1R-(1 α ,2 α ,4 β)]-(1-Aminomethyl-2,4-dimethyl-cyclopentyl)-acetic acid;
[1S-(1 α ,2 α ,4 β)]-(1-Aminomethyl-2,4-dimethyl-cyclopentyl)-acetic acid;
15 [1R-(1 α ,2 β ,4 α)]-(1-Aminomethyl-2,4-dimethyl-cyclopentyl)-acetic acid;
[1S-(1 α ,2 β ,4 α)]-(1-Aminomethyl-2,4-dimethyl-cyclopentyl)-acetic acid;
[1R-(1 α ,2 β ,4 β)]-(1-Aminomethyl-2,4-dimethyl-cyclopentyl)-acetic
20 acid; and
[1S-(1 α ,2 β ,4 β)]-(1-Aminomethyl-2,4-dimethyl-cyclopentyl)-acetic acid.

6. A method according to Claim 1 wherein the compound administered is named (3S, 4S)-1-Aminomethyl-3,4-dimethyl-cyclopentyl)-acetic acid.
- 25 7. A method according to Claim 1 wherein the compound administered is named (3S, 5R)-3-Aminomethyl-5-methyl-octanoic acid.
8. A method according to Claim 1 wherein the sleep disorder is insomnia.

-35-

9. A method according to Claim 1 wherein the sleep disorder is insomnia, and the compound administered is named (3S, 4S)-(1-Aminomethyl-3,4-dimethyl-cyclopentyl)-acetic acid.
10. A method according to Claim 1 wherein the sleep disorder is insomnia, and the compound administered is named (3S, 5R)-3-Aminomethyl-5-methyl-octanoic acid.